

# Der nicht tastbare Hoden und dann?

## Praktische Tipps für eine eher seltene klinische Situation

Anne-Karoline Ebert

### Literatur

1. Zakaria OM et al. Examination under anesthesia for management of impalpable undescended testis: a traditional technique revisited. *World J Surg* 2013; 37(5): 1125–9
2. Radmayr et al. EAU Guidelines. Edn. presented at the EAU Annual Congress Amsterdam 2022. ISBN 978-94-92671-16-5.
3. Berger C et al. Nonpalpable testes: Ultrasound and contralateral testicular hypertrophy predict the surgical access, avoiding unnecessary laparoscopy. *J Pediatr Urol* 2018; 14(2): 163.e1-163.e7
4. Tasian GE und Copp HL. Diagnostic performance of ultrasound in nonpalpable cryptorchidism: a systematic review and meta-analysis. *Pediatrics* 2011; 127(1): 119–28
5. Hodhod A et al. Testicular hypertrophy as a predictor for contralateral monorchism: Retrospective review of prospectively recorded data. *J Pediatr Urol* 2016; 12(1): 34.e1-5
6. Mah LW et al. Non-palpable testis: is management consistent and objective? *J Pediatr Urol* 2020; 16(1): 62–8
7. Sturm R et al. Blind ending vessels on diagnostic laparoscopy for nonpalpable testis: Is a nubbin present? *J Pediatr Urol* 2017; 13(4): 392.e1-392.e6
8. Nataraja RM et al. Is routine excision of testicular remnants in testicular regression syndrome indicated? *J Pediatr Urol* 2015; 11(3): 151.e1-5
9. Mentessidou A et al. Laparoscopic versus open orchiopexy for palpable undescended testes: Systematic review and meta-analysis. *J Pediatr Surg* 2022; 57(4): 770–5
10. Yu C et al. Evaluation of Fowler-Stephens orchiopexy for high-level intra-abdominal cryptorchidism: A systematic review and meta-analysis. *Int J Surg* 2018; 60: 74–87
11. Braga LH et al. Gubernaculum Testis and Cremasteric Vessel Preservation during Laparoscopic Orchiopexy for Intra-Abdominal Testes: Effect on Testicular Atrophy Rates. *J Urol* 2019; 201(2): 378–85