

# Urolithiasis: Hat die Extrakorporale Stoßwellentherapie noch eine Chance?

Konrad Wilhelm

## Literatur

1. Heers H et al. Urolithiasis in Germany: Trends from the National DRG Database. Urol Int 2022; 108(6): 589–95
2. Seitz C et al. „S2k-Leitlinie zur Diagnostik, Therapie und Metaphylaxe der Urolithiasis“, AWMF-Registernummer: 043–025. ([https://www.awmf.org/uploads/tx\\_szleitlinien/043-025l\\_S2k\\_Diagnostik\\_Therapie\\_Metaphylaxe\\_Urolithiasis\\_2019-07\\_1.pdf](https://www.awmf.org/uploads/tx_szleitlinien/043-025l_S2k_Diagnostik_Therapie_Metaphylaxe_Urolithiasis_2019-07_1.pdf)) Zugegriffen: 15.02.2021
3. Klein J et al. Extrakorporale Stoßwellenlithotripsie. Urologe A 2018; 57(4): 463–73
4. Schuler TD et al. Medical expulsive therapy as an adjunct to improve shockwave lithotripsy outcomes: a systematic review and meta-analysis. J Endourol 2009; 23(3): 387–93
5. Liu LR et al. Percussion, diuresis, and inversion therapy for the passage of lower pole kidney stones following shock wave lithotripsy. Cochrane Database Syst Rev 2013; (12): CD008569
6. Schoenthaler M et al. Feasibility of an Updated Randomised Controlled Trial on Surgical Urolithiasis Treatments: The Pilot Trial for the German Endoscopic versus Shock Wave Therapy Study (GESS). Eur Urol Focus 2022; 8(1): 271–5